

# Uniform Lyndon Interpolation for $\mathbf{N}^+ \mathbf{A}_{m,n}$

---

Yuta Sato

18th Asian Logic Conference in Kyoto Sangyo Univ.

September 12th, 2025

Kobe University, Japan

The slides are available online at:


[cannorin.net/math/alc2025.pdf](http://cannorin.net/math/alc2025.pdf)



(will be displayed again at the end)

# Bonus: the Kripke game!

Daily Challenge: 00:33:18 until the next game.



Guess! (♡1)

Enter modal formula

Check! (♡1)

YOU WIN!



$\Diamond(\Box p \rightarrow p)$

$\Box p \rightarrow p$

$\Diamond\Box\perp$



I made a Wordle-like game  
where you guess the shape of a  
Kripke frame, just with formulas.  
Give it a try!

[cannorin.net/kripke](https://cannorin.net/kripke)



I proved that the logic  $\mathbf{N}^+ \mathbf{A}_{m,n}$  enjoys  
Uniform Lyndon interpolation property,  
with a new method called propositionalization.

This talk is based on:

Yuta Sato. Uniform Lyndon interpolation for the pure logic of necessitation with a modal reduction principle. *Journal of Logic and Computation*, to appear. [arXiv:2503.10176](https://arxiv.org/abs/2503.10176).

# Table of contents

The Logic  $\mathbf{N}^+ \mathbf{A}_{m,n}$

Uniform Lyndon Interpolation Property

The Propositionalization Method

The Main Theorem

# The Logic $N^+A_{m,n}$

---

# What is $\mathbf{N}^+ \mathbf{A}_{m,n}$ ?

$$\mathbf{N} := \mathbf{Cl} + \frac{\varphi}{\Box \varphi}$$

$$\mathbf{N}^+ \mathbf{A}_{m,n} := \mathbf{N} + \frac{\frac{\neg \Box \varphi}{\neg \Box \Box \varphi}}{} + \Box^n \varphi \rightarrow \Box^m \varphi$$

- $\mathbf{Cl}$ : the classical propositional logic
- $\mathbf{N}$ : the pure logic of necessitation (Fitting et al. 1992)
  - also obtained from the logic  $\mathbf{K}$  by removing its  $\mathbf{K}$  axiom
- $\frac{\neg \Box \varphi}{\neg \Box \Box \varphi}$ : required by the semantics\*
- $\Box^n \varphi \rightarrow \Box^m \varphi$ : a generalized reflexivity/transitivity axiom

---

\*the completeness does not hold without it. no deep dive today

# $N^+A_{m,n}$ vs. normal modal logics

## Fact (Kurahashi and S.)

$$N^+A_{m,n} \subsetneq K + \Box^n \varphi \rightarrow \Box^m \varphi$$

## Proof.

The rule  $\frac{\neg \Box \varphi}{\neg \Box \Box \varphi}$  is admissible in  $K$ . The rest is trivial.  $\square$

## Fact (Kurahashi and S.)

$N^+A_{m,n}$  has the finite frame property (ffp) for every  $m, n \in \mathbb{N}$

It is still unknown to this day whether  $K + \Box^n \varphi \rightarrow \Box^m \varphi$  has ffp

➡ The lack of the  $K$  axiom is indeed a massive difference



# The sequent calculus for $N^+A_{m,n}$

A sequent calculus  $G_{N^+A_{m,n}}$  is obtained from **LK** by adding:

$$\frac{\Rightarrow \varphi}{\Rightarrow \Box \varphi} \text{ (nec)}$$

$$\frac{\Box \varphi \Rightarrow}{\Box \Box \varphi \Rightarrow} \text{ (rosbox, when } m = 0 \text{ and } n \geq 2)$$

$$\frac{\Box^m \varphi, \Box^n \varphi, \Gamma \Rightarrow \Delta}{\Box^n \varphi, \Gamma \Rightarrow \Delta} \text{ (accL, when } n > m)$$

$$\frac{\Gamma \Rightarrow \Delta, \Box^m \varphi, \Box^n \varphi}{\Gamma \Rightarrow \Delta, \Box^m \varphi} \text{ (accR, when } m > n)$$

## Proposition (S.)

- $G_{N^+A_{m,n}}$  proves  $\Gamma \Rightarrow \Delta$  iff  $N^+A_{m,n}$  proves  $\bigwedge \Gamma \rightarrow \bigvee \Delta$
- $G_{N^+A_{m,n}}$  admits cut elimination

# Uniform Lyndon Interpolation Property

---

## CIP and LIP (1/2)

Let  $V^+(\varphi)$  and  $V^-(\varphi)$  denote the set of variables that occur in  $\varphi$  positively and negatively, resp. Let also  $V(\varphi) = V^+(\varphi) \cup V^-(\varphi)$

$L$  is said to enjoy Craig interpolation property (CIP) if for every  $\varphi, \psi$  s.t.  $L \vdash \varphi \rightarrow \psi$ , there is  $\chi$  s.t.:

1.  $V(\chi) \subseteq V(\varphi) \cap V(\psi)$ ;
2.  $L \vdash \varphi \rightarrow \chi$  and  $L \vdash \chi \rightarrow \psi$ .

Such  $\chi$  is called an interpolant of  $\varphi \rightarrow \psi$

## CIP and LIP (2/2)

Let  $V^+(\varphi)$  and  $V^-(\varphi)$  denote the set of variables that occur in  $\varphi$  positively and negatively, resp. Let also  $V(\varphi) = V^+(\varphi) \cup V^-(\varphi)$

$L$  is said to enjoy Lyndon interpolation property (LIP) if for every  $\varphi, \psi$  s.t.  $L \vdash \varphi \rightarrow \psi$ , there is  $\chi$  s.t.:

1.  $V^\bullet(\chi) \subseteq V^\bullet(\varphi) \cap V^\bullet(\psi)$  ( $\bullet \in \{+, -\}$ );
2.  $L \vdash \varphi \rightarrow \chi$  and  $L \vdash \chi \rightarrow \psi$ .

Such  $\chi$  is called an interpolant of  $\varphi \rightarrow \psi$

## UIP and ULIP (1/2)

$L$  is said to enjoy Uniform interpolation property (UIP) if for any  $\varphi$  and any finite set of variables  $P$ , there is  $\chi$  s.t.

1.  $V(\chi) \subseteq V(\varphi) \setminus P$ ;
2.  $L \vdash \varphi \rightarrow \chi$ ;
3.  $L \vdash \chi \rightarrow \psi$  for any  $\psi$  s.t.  $L \vdash \varphi \rightarrow \psi$  and  $V(\psi) \cap P = \emptyset$ .

Such  $\chi$  is called a post-interpolant of  $(\varphi, P)$

## UIP and ULIP (2/2)

$L$  is said to enjoy Uniform Lyndon interpolation property (ULIP) if for any  $\varphi$  and any finite sets of variables  $P^+, P^-$ , there is  $\chi$  s.t.

1.  $V^\bullet(\chi) \subseteq V^\bullet(\varphi) \setminus P^\bullet$  ( $\bullet \in \{+, -\}$ );
2.  $L \vdash \varphi \rightarrow \chi$ ;
3.  $L \vdash \chi \rightarrow \psi$  for any  $\psi$  s.t.  $L \vdash \varphi \rightarrow \psi$  and  $V^\bullet(\psi) \cap P^\bullet = \emptyset$  ( $\bullet \in \{+, -\}$ ).

Such  $\chi$  is called a post-interpolant of  $(\varphi, P^+, P^-)$

## Several facts on the interpolation properties (1/2)

### Fact

- If  $L$  has UIP, then  $L$  has CIP
- If  $L$  has LIP, then  $L$  has CIP
- If  $L$  has ULIP, then  $L$  has both UIP and LIP (Kurahashi 2020)

### Fact (Kurahashi 2020)

- The classical propositional logic **CI** enjoys ULIP
- The modal logic **K** enjoys ULIP

## Several facts on the interpolation properties (2/2)

The situation is complicated for the extensions of **K**:

### Fact

- **KT** = **K** +  $\Box\varphi \rightarrow \varphi$  enjoys ULIP (Kurahashi 2020)
- For  $m > 0$ , **K** +  $\Box\varphi \rightarrow \Box^m\varphi$  enjoys CIP (Gabbay 1972) and LIP (Kuznets 2016)
- **K4** = **K** +  $\Box\varphi \rightarrow \Box\Box\varphi$  lacks UIP (Bílková 2007)
- **K** +  $\Box\Box\varphi \rightarrow \Box\varphi$  lacks even CIP (Marx 1995)

**K** +  $\Box^n\varphi \rightarrow \Box^m\varphi$ , in general, may or may not enjoy them

➔ What happens if we weaken it to  $\mathbf{N}^+\mathbf{A}_{m,n}$ ?



# The Propositionalization Method

---

## Propositionalization, in short

ULIP of a logic is sometimes proven by embedding it to some weaker logic where ULIP is already known:

### Example

Through the boxdot translation, ULIP of **K** implies ULIP of **KT**, and the failure of it in **S4** implies that of **K4**

I gave a sufficient condition on such embeddings:

### Theorem (S.)

For any logics  $L \subseteq M$ , if there is a translation with certain properties, propositionalization, of  $M$  into  $L$ , and  $L$  has ULIP, then so does  $M$

## Propositionalization, in detail (1/3)

Given a logic  $X$ , let  $\mathcal{L}_X$  designate the language of  $X$ .

Consider logics  $L$  and  $M$  s.t.  $\mathcal{L}_L \subseteq \mathcal{L}_M$  and  $L \subseteq M$ .

Now we want to *propositionalize* any  $\mathcal{L}_M$ -formula that is not expressible in  $\mathcal{L}_L$ :

### Definition

Let  $L'$  be the same logic as  $L$ , but its propositional variables extended by adding a fresh one  $p_\varphi$  for every  $\varphi \in \mathcal{L}_M$ .

### Definition

Let  $\sigma : \mathcal{L}'_L \rightarrow \mathcal{L}_M$  be the substitution that replaces every  $p_\varphi$  with  $\varphi$ , then  $L' \vdash \rho$  implies  $M \vdash \sigma(\rho)$  for any  $\rho \in \mathcal{L}'_L$ .

## Propositionalization, in detail (2/3)

### Definition

A pair of translations  $\sharp, \flat : \mathcal{L}_M \rightarrow \mathcal{L}'_L$  is called a propositionalization of  $M$  into  $L$  if the following are met:

(Embeddable)  $M \vdash \varphi \rightarrow \psi$  implies  $L' \vdash \varphi^\flat \rightarrow \psi^\sharp$ ;

(Invertible)  $M \vdash \sigma(\varphi^\sharp) \rightarrow \varphi$  and  $M \vdash \varphi \rightarrow \sigma(\varphi^\flat)$ ;

(Polarity-preserving) For  $(\bullet, \circ) \in \{(+, -), (-, +)\}$ ,  $\natural \in \{\sharp, \flat\}$ :

- $p \in V^\bullet(\varphi^\natural)$  implies  $p \in V^\bullet(\varphi)$ ;
- $p_\psi \in V^\bullet(\varphi^\natural)$  implies  $V^\bullet(\psi) \subseteq V^\bullet(\varphi)$ ,  $V^\circ(\psi) \subseteq V^\circ(\varphi)$ .

## Propositionalization, in detail (3/3)

### Theorem (S.)

If there is a propositionalization  $(\sharp, \flat)$  of  $M$  into  $L$ , and  $L$  has ULIP, then  $M$  does also

### Proof (outline).

Take any  $\varphi$ ,  $P^+$ ,  $P^-$ . We extend  $P^\bullet$  to  $Q^\bullet$  by adding every  $p_\psi \in V(\varphi^\flat)$  s.t.  $V^\bullet(\psi) \cap P^\bullet \neq \emptyset$ . By ULIP of  $L$ , we get a post-interpolant  $\chi'$  of  $(\varphi^\flat, Q^+, Q^-)$ . Then, embeddability, invertibility, and polarity-preservingness of  $\sharp, \flat$  assert that  $\chi = \sigma(\chi')$  is indeed a post-interpolant of  $(\varphi, P^+, P^-)$  in  $M$ .  $\square$

# The Main Theorem

---

# The Main Theorem

## Theorem (S.)

There is a propositionalization  $(\sharp, \flat)$  of  $\mathbf{N}^+ \mathbf{A}_{m,n}$  into  $\mathbf{CI}$

## Proof (outline).

We construct such  $\sharp, \flat$  that each additional rule in  $\mathbf{G}_{\mathbf{N}^+ \mathbf{A}_{m,n}}$  can be *emulated* in  $\mathbf{LK}$ :  $\mathbf{G}_{\mathbf{N}^+ \mathbf{A}_{m,n}} \vdash \Gamma \Rightarrow \Delta$  implies  $\mathbf{LK} \vdash \Gamma^\flat \Rightarrow \Delta^\sharp$ , then embeddability naturally holds. We also ensure invertibility and polarity-preservingness by adding just the right amount of information to enable such emulation.  $\square$

## Corollary

$\mathbf{N}^+ \mathbf{A}_{m,n}$  enjoys ULIP!

## Summing it up (1/2)

It is known that  $\mathbf{K} + \Box^n \varphi \rightarrow \Box^m \varphi$  does not, in general, enjoy all of CIP, LIP, UIP, and ULIP:

- $\mathbf{K4} = \mathbf{K} + \Box \varphi \rightarrow \Box \Box \varphi$  lacks UIP
- $\mathbf{K} + \Box \Box \varphi \rightarrow \Box \varphi$  lacks even CIP

However,  $\mathbf{N}^+ \mathbf{A}_{m,n}$  enjoy all of them for every  $m, n \in \mathbb{N}$ !



## Summing it up (1/2)

It is known that  $\mathbf{K} + \Box^n \varphi \rightarrow \Box^m \varphi$  does not, in general, enjoy all of CIP, LIP, UIP, and ULIP:

- $\mathbf{K4} = \mathbf{K} + \Box \varphi \rightarrow \Box \Box \varphi$  lacks UIP
- $\mathbf{K} + \Box \Box \varphi \rightarrow \Box \varphi$  lacks even CIP

However,  $\mathbf{N}^+ \mathbf{A}_{m,n}$  enjoy all of them for every  $m, n \in \mathbb{N}$ !

### Open Problem

To what extent the presence of the  $\mathbf{K}$  axiom is *harmful* for a logic in terms of interpolation properties?

- Is there a logic between  $\mathbf{N4}$  and  $\mathbf{K4}$  that lacks UIP?
- Is there a logic between  $\mathbf{N} + \Box \Box \varphi \rightarrow \Box \varphi$  and  $\mathbf{K} + \Box \Box \varphi \rightarrow \Box \varphi$  that lacks CIP?

## Summing it up (2/2)

We also developed a general method for proving ULIP:

### **Theorem (S.)**

For any logics  $L \subseteq M$ , if there is a propositionalization of  $M$  into  $L$ , and  $L$  has ULIP, then so does  $M$

## Summing it up (2/2)

We also developed a general method for proving ULIP:

### Theorem (S.)

For any logics  $L \subseteq M$ , if there is a propositionalization of  $M$  into  $L$ , and  $L$  has ULIP, then so does  $M$

### Open Problems

- Can we possibly say that if ULIP holds, then some nontrivial propositionalization exists? For example, can we construct propositionalizations of **K** into **N** or **CI**?
- Can we characterize a syntactic property on sequent calculi that corresponds to the existence of a propositionalization? (e.g. lemhoff 2019, Akbar Tabatabai & Jalali 2025)

# Thanks!

That's all!

The Slides



[cannorin.net/math/alc2025.pdf](http://cannorin.net/math/alc2025.pdf)

The Kripke Game



[cannorin.net/kripke](http://cannorin.net/kripke)

## **Appendix & References**

---

# References

This talk is based on the paper indicated by ★:

- ★ Yuta Sato. Uniform Lyndon interpolation for the pure logic of necessitation with a modal reduction principle. *Journal of Logic and Computation*, to appear. *arXiv:2503.10176*.
- Melvin C. Fitting, V. Wiktor Marek, and Mirosław Truszczyński. The pure logic of necessitation. *Journal of Logic and Computation*, 2(3):349-373, 1992.
- Taishi Kurahashi. The provability logic of all provability predicates. *Journal of Logic and Computation*, 34(6):1108-1135, 2024.
- Taishi Kurahashi and Yuta Sato. The Finite Frame Property of Some Extensions of the Pure Logic of Necessitation. *Studia Logica*, to appear.
- Taishi Kurahashi. Uniform Lyndon interpolation property in propositional modal logics. *Archive for Mathematical Logic*, 59(5-6):659-678, 2020.

# References

- Dov M. Gabbay. Craig's interpolation theorem for modal logics. *Conference in Mathematical Logic — London '70*, 111–127, 1972.
- Roman Kuznets. Proving Craig and Lyndon Interpolation Using Labelled Sequent Calculi. *Logics in Artificial Intelligence*, 320–335, 2016.
- Marta Bílková. Uniform Interpolation and Propositional Quantifiers in Modal Logics. *Studia Logica*, 85(1):1-31, 2007.
- Maarten Marx. Algebraic Relativization and Arrow Logic. ILLC Dissertation Series. Institute for Logic, Language and Computation, 1995.
- Rosalie Iemhoff. Uniform interpolation and the existence of sequent calculi. *Annals of Pure and Applied Logic*, 170(11):102711, 2019.
- A. Akbar Tabatabai, R. Jalali, Universal proof theory: Semi-analytic rules and Craig interpolation, *Annals of Pure and Applied Logic*, 176(1):103509, 2025.